



Don't Ban Chlorine—Use It Wisely

Recently, various activist and environmental groups have called for a ban on the use of chlorine and chlorine-containing compounds (1,2). The questions at issue are 1) is it possible, and 2) is it wise? Of course, chlorine, as chloride ion, is an essential component of the body, where it occurs at high levels in plasma and interstitial fluid. There probably is no controversy concerning this aspect of chlorine.


However, is it really possible to ban chlorine-containing compounds from the environment? Many do not realize the great number and ubiquitous presence of such substances in the environment (3,4). Burning of wood, a fuel with possible increased use, both for domestic and industrial purposes, leads to the release of methyl chloride, a weak alkylating agent (5). Marine organisms form chloroform, carbon tetrachloride, chlorine-containing haloforms, and longer-chain compounds containing various halogens. It was once estimated that the natural formation of carbon tetrachloride was greater than industrial production in the entire Northern Hemisphere (6). The comment might be made that these examples are all aliphatic compounds; that only the more persistent cyclic compounds containing chlorine should be banned. But nature, the consummate master of synthesis, has an appreciable number of cyclic organics which contain chlorine in the repertoire. Ochratoxin, made by the fungus *Aspergillus ochraceus* (7), is an example of a toxic material with chlorine on an aromatic ring. On the other hand, naturally occurring chlorinated compounds such as chloramphenicol, chlortetracycline, and griseofulvin have been used to advantage against bacterial or mycotic diseases of humans (8). Various chlorinated phenols with antibiotic properties have been isolated from naturally occurring organisms, especially marine algae (9,10). A survey of many marine organisms showed that numerous products with antiviral, antimicrobial, and antineoplastic properties were present (11).

Not only marine organisms synthesize chlorinated substances; lichens (12,13), and even some plants, seem capable of incorporating chlorine atoms into their metabolites (14). Furthermore, a substance with a structure akin to that of a chlorinated dibenzofuran has been isolated from soil under decomposed roots of a type of eucalyptus tree (15,16). Thus, nature furnishes chlorinated compounds similar to some of the hazardous materials presumed to result only from technological activity.

Is it wise to ban chlorine and chlorine-containing compounds? Part of the reason for the longer life span of our population is the availability, due to chlorination, of a water supply free of the organisms that cause typhoid, cholera, and other infectious diseases. It is often stated that other methods of disinfecting water could be used. True, but their disinfecting power usually does not extend beyond the actual disinfection period, while with chlorine there is a residual period when the action remains. With the increase in urban population of the United States and the resultant more crowded cities, the residuum of action, as from chlorine, is probably desirable.

Furthermore, to ban chlorinated compounds may limit discov-

eries of potential new therapeutic agents. For example, a trichlorodibromooctene isolated from the red marine algae *Portiaria hornemannii* had an unusually high cytotoxic effect on chemoresistant human brain, kidney, and colon tumor cell lines in culture, leading to selection by the drug development group of the National Cancer Institute for further study (17). But if chlorine compounds are banned, further development may be "illegal." Fortunately, an anti-chlorine mentality has not prevented the development of 2-chlorodeoxyadenosine for the treatment of hairy-cell leukemia.



1,3,6,8,11,13-Hexachloro-4,10-dihydroxydinaphtho[2,1-b',1',1'-d]furan-5,9-dione is naturally occurring.

Currently, thousands of otherwise healthy but breast cancer-prone women are enrolled in clinical studies which use tamoxifen as a prophylactic agent against breast cancer. The hepatocarcinogenicity of tamoxifen in rats (18) and its propensity to form numerous DNA adducts (19) casts some doubt on the wisdom of this effort. On the other hand, the chlorinated analogue of tamoxifen, toremifene, although it also suppressed the spontaneous endocrine and mammary tumors of rats, did not cause liver cancer and did not form DNA adducts (19). Accordingly, based on animal studies, the chlorinated analogue appears safer than tamoxifen.

Thus, chlorine and its compounds should not be banned but used safely and properly in situations where their unique properties are necessary. Instead of an outright ban, scientifically based risk-benefit evaluations should be made for each substance.

Elizabeth K. Weisburger
Scientist Emeritus

REFERENCES

- Hileman B. Concerns broaden over chlorine and chlorinated hydrocarbons. Chem Eng News April 19:11-20 (1993).
- Amato I. The crusade against chlorine. Science 261:152-154 (1993).
- Siuda JF, DeBernardis JF. Naturally occurring halogenated organic compounds. Lloydia 36:107-143 (1973).
- Hoyt SD, Rasmussen RA. Determining trace gases in air and seawater. Adv Chem Ser 209:31-56 (1985).
- Tassios S, Packham DR. The release of methylchloride from biomass burning in Australia. J Air Pollut Control Assoc 35:41-42 (1985).
- Lovelock JE. Natural halocarbons in the air and in the sea. Nature 256:193-194 (1975).
- van der Merwe, KF, Steyn PS, Fourie L, Scott DB, Theron JJ. Ochratoxin A, a toxic metabolite produced by *Aspergillus ochraceus* Wilh. Nature 205:1112-1113 (1965).
- IARC. IARC monographs on the evaluation of the carcinogenic risk of chemicals to man, vol 10. Some naturally occurring substances. Lyon:International Agency for Research on Cancer, 1976.
- Kavanagh F, Hervey A, Robbins WJ. Antibiotic substances from Basidiomycetes. IX. *Drosophila subvariate* (Batsch ex Fr.) Quel. Proc Natl Acad Sci USA 38:555-560 (1952).
- Weisburger EK. Halogenated substances: environmental and industrial materials. In: Toxicology of halogenated hydrocarbons (Khan MAQ, Stanton RH, eds). New York:Pergamon Press, 1981; 3-21.
- Rinehart KL Jr. Shaw PD, Shield LS, Gloer JB, Harbour GC, Koker MES,

- et al. Marine natural products as sources of antiviral, antimicrobial, and antineoplastic agents. *Pure Appl Chem* 53:795-817 (1981).
12. Miller MW. The Pfizer handbook of microbial metabolites. New York: McGraw-Hill, 1961.
 13. Huneck S, Sundholm G, Follman NG. 3-Chordivaricatsaure, ein neues Depsid aus *Thelasma*-arten. *Phytochemistry* 19:645-649 (1980).
 14. Segall HF, Krick TP. Pyrrolizidine alkaloids: organohalogen derivative isolated from *Senecio jacobaea* (tansy ragwort). *Toxicol Lett* 4:193-198 (1980).
 15. Cameron DW, Sidell MD. 1,3,6,8,11,13-Hexachloro-4,10-dihydroxydinaphtho [2.1-b:1',1'-d] furan-5,9-dione. A polychloro quinone from green soils. *Aust J Chem* 31:1323-1333 (1978).
 16. Cameron DW, Feutrill GI, Pannan LJH. Synthesis of a natural polychloro dinaphthofuran quinone. *Tetrahedron Lett* 21:1385-1386 (1980).
 17. Fuller RW, Cardellina JH II, Kato Y, Brinen LS, Clardy J, Snader KM, Boyd MR. A pentahalogenated monoterpene from the red algae *Portieria hornemannii* produces a novel, cytotoxicity profile against a diverse panel of human tumor cell lines. *J Med Chem* 35:3007-3011 (1992).
 18. Greaves P, Goonetilleke R, Nunn G, Topham J, Orton T. Two-year carcinogenicity study of tamoxifen in Alderley Park Wistar-derived rats. *Cancer Res* 53:3919-3924 (1993).
 19. Hard GC, Iatropoulos MJ, Jordan K, Radi L, Kaltenberg, OP, Imondi AR, Williams GM. Major difference in the hepatocarcinogenicity and DNA adduct forming ability between toremifene and tamoxifen in female Crl:CD(BR) rats. *Cancer Res* 53:4534-4541 (1993).

ASBMB Fall Symposia Series

Genetic and Biochemical Approaches for Studying Cell Death

Organizer: Stanley J. Korsmeyer

October 7-10, 1994

Granlibakken, Lake Tahoe, California

Mechanisms of Regulated Intracellular Protein Degradation

Organizers: Stuart M. Arfin and Ralph A. Bradshaw

October 14-17, 1994

Whistler, British Columbia, Canada

Oligonucleotide Selection and Molecular Diversity

Organizers: Jack D. Keene and David L. Bartel

October 28-31, 1994

Granlibakken, Lake Tahoe, California

For further information call:

ASBMB Society Office

9650 Rockville Pike

Bethesda, MD 20814-3992

Phone: 301-530-7145 or FAX: 301-571-1824